

EXXON RESEARCH AND ENGINEERING COMPANY

P.O. BOX 101, FLORHAM PARK, NEW JERSEY 07932

M. B. GLASER
Manager
Environmental Affairs Programs

Cable: ENGREXXON, N.Y.

November 12, 1982

CO₂ "Greenhouse" Effect


82EAP 266

TO: See Distribution List Attached

Attached for your information and guidance is briefing material on the CO₂ "Greenhouse" Effect which is receiving increased attention in both the scientific and popular press as an emerging environmental issue. A brief summary is provided along with a more detailed technical review prepared by CPPD.

The material has been given wide circulation to Exxon management and is intended to familiarize Exxon personnel with the subject. It may be used as a basis for discussing the issue with outsiders as may be appropriate. However, it should be restricted to Exxon personnel and not distributed externally.

Very truly yours,



M. B. GLASER

MBG:rva

Attachments

H. N. WEINBERG

NOV 15 1982



Distribution List:

R. W. Cohen
M. J. Connor, Jr.
E. E. David, Jr.
C. M. Eidt, Jr.
W. R. Epperly
R. L. Hirsch
T. G. Kaufmann
D. G. Levine
G. H. Long
J. R. Riley
H. R. Savage
A. Schriesheim
J. F. Taylor
D. T. Wade
H. N. Weinberg

SUMMARY

Atmospheric monitoring programs show the level of carbon dioxide in the atmosphere has increased about 8% over the last twenty-five years and now stands at about 340 ppm. This observed increase is believed to be the continuation of a trend which began in the middle of the last century with the start of the Industrial Revolution. Fossil fuel combustion and the clearing of virgin forests (deforestation) are believed to be the primary anthropogenic contributors although the relative contribution of each is uncertain.

The carbon dioxide content of the atmosphere is of concern since it can affect global climate. Carbon dioxide and other trace gases contained in the atmosphere such as water vapor, ozone, methane, carbon monoxide, oxides of nitrogen, etc. absorb part of the infrared rays reradiated by the earth. This increase in absorbed energy warms the atmosphere inducing warming at the earth's surface. This phenomenon is referred to as the "greenhouse effect".

Predictions of the climatological impact of a carbon dioxide induced "greenhouse effect" draw upon various mathematical models to gauge the temperature increase. The scientific community generally discusses the impact in terms of doubling of the current carbon dioxide content in order to get beyond the noise level of the data. We estimate doubling could occur around the year 2090 based upon fossil fuel requirements projected in Exxon's long range energy outlook. The question of which predictions and which models best simulate a carbon dioxide induced climate change is still being debated by the scientific community. Our best estimate is that doubling of the current concentration could increase average global temperature by about 1.3° to 3.1° C. The increase would not be uniform over the earth's surface with the polar caps likely to see temperature increases on the order of 10° C and the equator little, if any, increase.

Considerable uncertainty also surrounds the possible impact on society of such a warming trend, should it occur. At the low end of the predicted temperature range there could be some impact on agricultural growth and rainfall patterns which could be beneficial in some regions and detrimental in others. At the high end, some scientists suggest there could be considerable adverse impact including the flooding of some coastal land masses as a result of a rise in sea level due to melting of the Antarctic ice sheet. Such an effect would not take place until centuries after a 3° C global average temperature increase actually occurred.

There is currently no unambiguous scientific evidence that the earth is warming. If the earth is on a warming trend, we're not likely to detect it before 1995. This is about the earliest projection of when the temperature